


Lecture 5b

Drugs that Inhibit RNA and Protein Synthesis, Gene Editing


Drugs as inhibitors of Transcription:

Initiation



RNA polymerase


Elongation



Fidaxomicin, Myxopyronin

- Transcription is the process of copying DNA to RNA
- RNA polymerase binds to a specific DNA sequence called the promoter
- RNAP generates a complementary copy of one strand, T is replaced by U.
- Termination occurs due to specific signals in the RNA sequence.

Termination



Bicyclomycin

Protein Synthesis – tRNA & Ribosomes

Role of different Ribosomal subunits

30S (Small) – RBS & mRNA codon/anticodon

50S (Large) – Peptide bond synthesis

Exit tunnel – new protein emerges

tRNA sites:

A – aminoacyl – next tRNA-AA binds

P – 1st tRNA-Met & growing peptide

E – empty tRNA leave from here

Initiation:

1. Ribosome binding site & rRNA interaction (Proks)/AUG scanning (Euks).

2. fMet-tRNA (Proks) or Met (Euks) in P site

Elongation:

1. New AA-tRNA in A site

2. Peptide bond formation (amino acid in A site added to C-term of peptide in P site)


3. Translocation (tRNA-peptide moves to P site)

4. tRNA exits

Termination:


1. Stop codon at A site

2. Termination factor (protein) adds water to cleave peptide from last tRNA




tRNAs

- The adapter molecules are called transfer RNAs or tRNAs.
- Contain:
 - a CCA sequence at 3' end where the amino acid is attached
 - a triplet anticodon to form base pairs with the appropriate mRNA codon
- The tRNA-AA is called a charged tRNA



Antibiotics that Inhibit Protein Synthesis



Genome Editing


Genome Editing – CRISPR Cas9

A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity

17 AUGUST 2012 VOL 337 SCIENCE www.sciencemag.org

Martin Jinek,^{1,2*} Krzysztof Chylinski,^{3,4*} Ines Fonfara,⁴ Michael Hauer,^{2†}
Jennifer A. Doudna,^{1,2,5,6‡} Emmanuelle Charpentier^{4‡}

Clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated (Cas) systems provide bacteria and archaea with adaptive immunity against viruses and plasmids by using CRISPR RNAs (crRNAs) to guide the silencing of invading nucleic acids. We show here that in a subset of these systems, the mature crRNA that is base-paired to trans-activating crRNA (tracrRNA) forms a two-RNA structure that directs the CRISPR-associated protein Cas9 to introduce double-stranded DNA breaks at target DNA. At sites complementary to the crRNA-guide sequence, the Cas9 HNH nuclease domain cleaves the complementary strand, whereas the Cas9 RuvC-like domain cleaves the non-complementary strand. The dual-tracrRNA:crRNA, when engineered as a single RNA chimeric molecule, directs sequence-specific Cas9 dsDNA cleavage. This family of endonucleases, which use dual-RNAs for target recognition, has the potential to exploit the Cas9 system for RNA-guided genome engineering.



The Nobel Prize in Chemistry
2020



© Nobel Prize Outreach. Photo:
Bernhard Ludewig
Emmanuelle
Charpentier
Prize share: 1/2




© Nobel Prize Outreach. Photo:
Brittany Hosea-Small
Jennifer A. Doudna
Prize share: 1/2

The Nobel Prize in Chemistry 2020 was awarded jointly to Emmanuelle Charpentier and Jennifer A. Doudna "for the development of a method for genome editing"

Key Concepts in Genome Editing

Repair of a targeted double strand break → modification of the genome at a defined location.



Original Sequence
--ATG.....GGGTGG**GCC**GATT...CGATAA--
--Met.....GlyTrpProIle...Arg



Deletion of one base
--ATG.....GGGTG**CC**GATT...CGATAA--
--Met.....GlyCysArgLeu.ArgIle...

Indels:

- $+/- 3n$ = addition or loss of amino acids
- $+/- 1$ or 2 bases = Frame shift. changes the amino acid sequence after the indel. The ribosome considers three bases (codon) relative to the start codon.

Key Concepts in Genome Editing

Repair method II – replacement of a segment of DNA



How to Cut at a Defined Location - Cas9 + Guide RNA

Cas9-RNA complex

- Cas9 – nuclease that cuts DNA after activation
- Guide RNA:
 - 5' end complementary to target sequence
 - 3' end required for Cas9 activity (tracrRNA)

Double stranded break
(3 bases from PAM)


2. After PAM recognition by Cas9, guide RNA unwinds DNA, by pairing with one DNA strand.

3. Cas9 cleaves both strands near site, generating a double strand break.

4. Double stranded break triggers DNA repair:

- Generation of indel
- Using injected replacement DNA for homologous repair


1. Cas9 Binds to PAM,




Altering the Genome Sequence with Cas9-CRISPR

Components to microinject:

1. Cas9 enzyme (nuclease)
2. Guide RNA, specific for site of cleavage, bound to the Cas9 protein
3. Copy of replacement DNA sequence (dsDNA)



1. Guide RNA directs Cas9 to desired site, by pairing with one DNA strand.
2. CRISPR cleaves both strands near site, generating a double strand break.
3. Double stranded break triggers DNA repair, using injected replacement DNA for homologous repair



(Video originally from Nature)

Also view:


<https://wyss.harvard.edu/media-post/gene-editing-mechanism-of-crispr-cas9/>

Using CRISPR-Cas9 to Correct Genetic Diseases

Human growth hormone (hGH)

Pituitary Dwarfism

(a) *GH1* codes for a pituitary growth hormone.



Between one in 14,000 and one in 27,000 babies born each year have some form of dwarfism.


(b) Normal versus GH1-deficient



1860 William Harrison and Charles Stratton - comedians and performers.

Components to microinject:

1. Cas9 enzyme (nuclease)
2. Guide RNA, specific for site of cleavage, bound to the Cas9 protein
3. Copy of replacement DNA sequence (dsDNA)



Human growth hormone gene

Possible PAM Sites

Wild type (normal)

R L E D G S P R T G Q I F K Q T Y S
-CTTGAGAGGCTGGAAAGATGGCAGCCCCCGGACTGGGCAGATCTCAAGCAGACCTACAGCAA-
-GAAACGTCTCCGACCTTCTACCGTCGGGGCCTGACCCGTCTAGAAGTTCGTCTGGATGTCGT-

Mutant (growth hormone non-functional)

R L E D G S P R T G Q N F K Q T Y S
-CTTGAGAGGCTGGAAAGATGGCAGCCCCCGGACTGGGCAGAACTCAAGCAGACCTACAGCAA-
-GAAACGTCTCCGACCTTCTACCGTCGGGGCCTGACCCGTCTAGAAGTTCGTCTGGATGTCGT-

CRISPR Repair of Growth Hormone Gene

>M13438.1:497-2129 Human growth hormone gene (HGH-N), complete cds
AGGATCCCAAGGCCAACCTCCCAGAACCACTCAGGGTCTGTGGACAGCTCACCTAGCTGCAATGGCTAC 70
AGGTAAGCGCCCTAAATCCCTTGGACAATGTGCTCTGAGGGAGAGGCAGCGACCTGTAGATGGGA 140
CGGGGGCACTAACCTCAGGGTTGGGTTCTGAATGTGAGTATGCCATCTAACCCCAGTATTGGCCA 210
ATCTCAGAAAGCTCCTGGCTCCCTGGAGGATGGAGAGAGAAAAACAAACAGCTCTGGAGCAGGGAGAGT 280
GTTGGCCTCTGCTCTCCGGCTCCCTCTGTTGCCCTCTGTTCTCCCAAGCTCCGGACGTCCCTGCT 350
CCTGGCTTTGGCCTGCTCTGCCCTGGCTCAAGAGGGCAGTGCCTCCCAACCATTCCCTATCC 420
AGGCTTTTGACAACGCTATGCTCCCGCCCATCGTCTGCACCAGCTGGCCTTGACACCTACAGGAGT 490
TTGTAAGCTCTGGGAATGGGTGCGCATCAGGGTGGCAGGAAGGGTGACTTCCCCCGTGGAAATA 560
AGAGGAGGAGACTAAGGAGCTCAGGGTTTCCCACCGCGAAAATGCAGGCAGATGAGCACACGCTGAG 630
CTAGGTTCCCAGAAAAGTAAATGGGAGCAGGTCTCAGCTCAGACCTGGTGGCGGTCTCTCTAGG 700
AAGAACGCTATATCCAAAGGAACAGAAAGTATTCAATTCTGCAGAACCCCCAGACCTCCCTGTTCTC 770
AGAGTCTATTCCGACACCCCTCAACAGGGAGGAAACACAACAGAAATCCGTGAGTGGATGCCCTCCCC 840
AGGCAGGGATGGGGAGACCTGTAGTCAGAGCCCCGGCAGCACAGCCAATGCCGTCTGCCCTGCG 910
AGAACCTAGAGCTGCTCCGCATCTCCCTGCTCATCCAGTCGTGGCTGGAGCCCGTGCAGTTCTCAG 980
GAGTGTCTCGCCAACAGCCTGGTGTACGGCGCTCTGACAGCAACGTCTATGACCTCTAAAGGACCTA 1050
GAGGAAGGCATCCAAACGCTGATGGGGTGAGGGTGGCGCCAGGGTCCCCAATCTGGAGCCCACTGAG 1120
CTTGAGAGACTGTGTTAGAGAAACACTGGCTGCCCTTTTAGCAGTCAGGCCCTGACCCAAGAGAAC 1190
TCACCTTATTCTCATTCCCTCGTAATCCTCCAGGCCCTTCTACACTGAAGGGAGGGAGGAAA 1260
TGAATGAATGAGAAAGGGAGGAAACAGTACCCAAAGCGCTGGCCTCTCCTCTTCACTTTGCAAG 1340
AGGCTGGAAGATGGCAGCCCCCGGACTGGGCAGATCTCAAGCAGACCTACAGCAAGTTGACACAAACT 1410
CACACAACGATGACGCACTACTCAAGAACTACGGGCTCTCTACTGCTTCAGGAAGGACATGGACAAGGT 1480
CGAGACATTCCCTGCGCATCGTCAGTGGCCTCTGTGGGGCAGCTGTGGCTCTAGCTGCCCGGGTGG 1550
CATCCCTGTGACCCCTCCCCAGTGCCTCTCTGGCCCTGCAAGTTGCCACTCCAGTGCACCCACCAGCCTG 1620
TCTTAATAAAATTAAAGTGCATCATT

Location of mutation
Isoleucine (I) to Asparagine (N)

- The cut site needs to be close to site of mutation so that the injected dsDNA repair template can be as short as possible.
- A **NGG** (PAM site) is needed for Cas9 to bind & then test whether the RNA is complementary to the DNA.
- There are four possible PAM sites in the DNA sequence on the bottom left. The PAM site closest to the mutation was selected so that the cut site is close to mutation site.

CRISPR Repair of Grown Hormone Gene

- The PAM site closest to the mutation was selected so that the cut site is close to mutation site.
- The targeting section of the guide RNA should have the same sequence as 5' to the XGG, 18 bases are required:
5'AGAUGGCAGCCCCCGGAC----- plus additional RNA needed for Cas9 function
- This RNA would cause cleavage of both the wild-type or mutant sequence since they are identical in this region. This is OK since the repair DNA will contain the wild-type sequence.
- The site of Cas9 cleavage is between the PAM and the guide RNA sequence.
- The injected DNA contains sequences on both sides of the ds break, causing the replacement of the sequences at the double stranded break due to repair.

Injected dsDNA for Homologous Repair

Possible PAM Sites

Wild type (normal)

DNA cuts by cas9

R L E D G S P R T G Q I F K Q T Y S
-CTTGAGAGGCTGGAAAGATGGCAGCCCCGGACTGGGCAGATCTTCAAGCAGACCTACAGCAA-
-AACGTCTCCGACCTTCTACCGTCGGGGCCTGACCCGTCTAGAAGTTCGTCTGGATGTCCTT-

Mutant (growth hormone non-functional)

R L E D G S P R T G Q N F K Q T Y S
-CTTGAGAGGCTGGAAAGATGGCAGCCCCGGACTGGGCAGAACTTCAAGCAGACCTACAGCAA-
-AACGTCTCCGACCTTCTACCGTCGGGGCCTGACCCGTCTTAGAAGTTCGTCTGGATGTCCTT-

CTTGAGAGGCTGGAAAGATGGCAGCCCCGGACTGGGCAGATCTTCAAGCAGACCTACAGCAA-GAACGTCTCCGACCTTCTACCGTCGGGGCCTGACCCGTCTAGAAGTTCGTCTGGATGTCCTT-

homology regions required for repair

Note that Cas9 will cut both the wild-type and the mutant, but repair will insert the wild-type sequence.

Mutant (growth hormone non-function)

-AGGCTGGAGATGGCAGCCCCGGACTGGCAGAACATTCAAGCAGACCTACAGCAA-
-TCCGACAGATGGCAGCCCCGGACTGGCAGAACATTCAAGCAGACCTACAGCAA-

guide RNA

Cas9

Editing Steps:

1. Cas9 binds to NGG (PAM)
2. Opens DNA if RNA is complementary to DNA
3. Cas9 cuts both strands
4. Double stranded break causes repair.
5. Injected template is used to repair, changing the DNA sequence between the two homologous regions.

